


[Web](#) [Images](#) [Video](#) [News](#) [Maps](#) [more »](#)

[Advanced Scholar Search](#)
[Scholar Preferences](#)
[Scholar Help](#)

Scholar [All articles](#) - [Recent articles](#) Results **1 - 10** of about **389** for **benzoyl isocyanate stationary phase polysaccharide**. (0.16 seconds)

[Enantiomeric Resolution of Chiral Sulfoxides on **Polysaccharide** Phases by HPLC](#) - [all 2 versions »](#)

SA MATLIN, ME TIRITAN - CHIRALITY, 1996 - doi.wiley.com

... last 10 years, derivatives of **polysaccharides** such as ... enantiomers and (S)- α -methyl-**benzyl isocyanate** were purchased ... 1. Structure of chiral **stationary phase**. ...

[Cited by 27](#) - [Related Articles](#) - [Web Search](#) - [BL Direct](#)

[Covalently bonded **polysaccharide** derivatives as chiral **stationary** phases in high-performance liquid ...](#) - [all 5 versions »](#)

P Franco, A Senso, L Oliveros, C Mingullón - Journal of Chromatography A, 2001 - Elsevier

... of selectors to be used in chiral **stationary** phases (CSPs). ... by reaction of the **polysaccharide** with a **benzoyl** chloride or a phenyl **isocyanate** in homogeneous ...

[Cited by 60](#) - [Related Articles](#) - [Web Search](#)

[... electrochromatography using polyacrylamide and **polysaccharide** derivatives as chiral **stationary** ...](#) - [all 4 versions »](#)

K Krause, M Girod, B Chankvetadze, G Blaschke - Journal of Chromatography A, 1999 - Elsevier

... as a better reproducibility of data and **stationary** beds ... 3) benzoin, (4) N-**benzoyl**-phenyl-glycin ... by reaction of the corresponding **isocyanate** with ...

[Cited by 86](#) - [Related Articles](#) - [Web Search](#)

[... of selected chiral sulfoxides using **polysaccharide**-type chiral **stationary** phases and polar organic, ...](#) - [all 5 versions »](#)

B Chankvetadze, C Yamamoto, Y Okamoto - Journal of Chromatography A, 2001 - Elsevier

... recognition ability among chiral **stationary** phases (CSPs ... benzylsulfinyl)benzoic acid **benzyl** ester (BSBABE ... 3,5-dichlorophenyl **isocyanate**, pyridine, tetrahydrofuran ...

[Cited by 44](#) - [Related Articles](#) - [Web Search](#)

[... , p-toluenesulfonylcarbamate, and benzoylcarbamates of cellulose and amylose as chiral **stationary** ...](#) - [all 3 versions »](#)

T Ikai, C Yamamoto, M Kamigaito, Y Okamoto - Chirality, 2005 - doi.wiley.com

... performance liquid chromatography; HPLC; chiral **stationary phase** ... used as chiral **stationary** phases (CSPs ... carbonate, thionyl chloride, **benzoyl isocyanate**, and 3,5 ...

[Cited by 6](#) - [Related Articles](#) - [Web Search](#) - [BL Direct](#)

[Polysaccharide derivatives as chiral **stationary** phases in HPLC](#) - [all 2 versions »](#)

Y Okamoto, Y Kaïda - J. High Resol. Chromatogr, 1990 - doi.wiley.com

... HPLC, development of a Chiral **Stationary Phase** (CSP) is ... on aralkylcarbamates such as **benzyl** and I ... of polysac- charides with corresponding **isocyanate** derivatives ...

[Cited by 13](#) - [Related Articles](#) - [Web Search](#)

[... characterisation and study of the resulting high-performance liquid chromatography chiral **stationary** ... - all 4 versions »](#)

A Senso, L Oliveros, C Minguillón - Journal of Chromatography A, 1999 - Elsevier

... 1. Preparation of the chiral **stationary** phases. ... suspended in pyridine and treated with an excess of the corresponding aryl **isocyanate** or **benzoyl** chloride (6 ...

[Cited by 29](#) - [Related Articles](#) - [Web Search](#)

[... \) benzamide Using Cellulose Tris \(3, 5-dichlorophenylcarbamate\) as a Chiral **Stationary Phase** - all 3 versions »](#)

B Chankvetadze, C Yamamoto, Y Okamoto - Chemistry Letters, 2000 - J-STAGE

... an excess of 3,5-dichlorophenyl **isocyanate** in dry ... Tris(3,5-dichlorophenylcarbamate) as a Chiral **Stationary Phase** ... of compound 1 containing a **benzyl** ester group ...

[Cited by 6](#) - [Related Articles](#) - [Web Search](#) - [BL Direct](#)

[Immobilized **Polysaccharide**-Based Chiral **Stationary** Phases for HPLC - all 3 versions »](#)

T Ikai, C Yamamoto, M Kamigaito, Y Okamoto - Polymer Journal, 2006 - J-STAGE

... 1 and in 1981, the chiral **stationary** phases (CSPs ... the same selector and an identical amount of diisocyanate. ... then heated in the presence of **benzoyl** peroxide as ...

[Cited by 10](#) - [Related Articles](#) - [Web Search](#) - [BL Direct](#)

[Impact of immobilized **polysaccharide** chiral **stationary** phases on enantiomeric separations - all 3 versions »](#)

I Ali, HY Aboul-Enein - J. Sep. Sci, 2006 - doi.wiley.com

... silica with tolylene-2,4-diisocyanate as a spacer ... 1-(3,4-di- methoxy-**benzoyl**)-6-quinolyl ...

However, immobilized **polysaccharide**-based chiral **stationary** phases have ...

[Cited by 6](#) - [Related Articles](#) - [Web Search](#) - [BL Direct](#)

Key authors: [Y Okamoto](#) - [B Chankvetadze](#) - [K Krause](#) - [C Yamamoto](#) - [A Senso](#)

Google

Result Page: [1](#) [2](#) [3](#) [4](#) [5](#) [6](#) [7](#) [8](#) [9](#) [10](#) [Next](#)

benzoyl isocyanate stationary phase

[Google Home](#) - [About Google](#) - [About Google Scholar](#)

©2008 Google